

WHAT IS CLAIMED IS:

1. A method for decontaminating contamination containing biological spores, comprising the steps of:

contacting the contamination with a spore germination composition comprising dipicolinic acid and calcium ions effective to cause germination of the spores; and,

applying a decontaminating solution to kill the germinated spores.
2. The method of claim 1, wherein the step of contacting the contamination with the spore germination composition effective to cause germination of the spores simultaneously with the step of applying a decontaminating solution to kill the germinated spores.
3. The method of claim 1, wherein the step of contacting the contamination with the spore germination composition effective to cause germination of the spores occurs prior to the step of applying a decontaminating solution to kill the germinated spores.
4. The method of claim 1, wherein the spore germination composition comprises from about 50 mM to about 90 mM dipicolinic acid.

5. The method of claim 1, wherein the calcium ions comprise calcium chloride.
6. The method of claim 5, wherein the spore germination composition comprises from about 50 mM to about 90 mM calcium chloride.
7. The method of claim 6, wherein the spore germination composition comprises from about 60 mM to about 80 mM calcium chloride.
8. The method of claim 1, wherein the spore germination composition comprises from about 0.8% w/w to about 5% w/w dipicolinic acid of the total spore germination composition.
9. The method of claim 1, wherein the spore germination composition further comprises water.
10. The method of claim 9, wherein the spore germination composition comprises from about 50% w/w to about 98% w/w water.
11. The method of claim 1, wherein the spore germination composition further comprises

a surfactant.

12. The method of claim 11, wherein the surfactant is selected from the group consisting of anionic surfactant and nonionic surfactant.
13. The method of claim 11, wherein the surfactant comprises at least one carbon chain of from about six carbon members or more.
14. The method of claim 11, wherein the surfactant comprises from about 5% w/w to about 15% w/w of the total spore germination composition.
15. The method of claim 1, wherein the decontaminating solution comprises enzymes.
16. The method of claim 1, wherein the decontaminating solution comprises a peroxygen compound.
17. A germination composition for decontaminating biological spores, comprising dipicolinic acid and calcium ions.

- 5
10
15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995
18. The germination composition of claim 17, wherein the calcium ions are supplied by calcium chloride.
19. A decontaminated surface made by the process comprising the steps of:
contacting a surface with a spore germination composition comprising dipicolinic acid and calcium ions effective to cause germination of the spores; and,
applying a decontaminating solution to kill the germinated spores.
20. A method for decontaminating a chemical-biological agent comprising the steps of:
solubilizing the chemical-biological agent with a microemulsion having a plurality of surfactants; and,
decontaminating the solubilized chemical-biological agent with a peracid.